

Comment on “Nongaussian Isocurvature Perturbations from Inflation”

In a recent Letter [1] a hybrid inflationary model with nongaussian density perturbations and a “blue” spectral index $n \geq 1$ was presented. The purpose of this comment is to point out that this model can be considered as a particular realization of the general framework for the spectrum and statistics of the CMBR we proposed in [2], based on the hypothesis of conformal invariance. The implication of this hypothesis is that density perturbations are nongaussian with a spectral index $n \geq 1$.

In [1] the spectrum of density fluctuations is generated by a free massive scalar field propagating in (approximately) de Sitter background. The main contribution to the energy density in this model comes from the modes with (comoving) wavelengths larger than the horizon scale H^{-1} . In addition, the mean value of the scalar field *vanishes* as a result of averaging over sufficiently large scales, and the density perturbations are quadratic in the scalar field fluctuations. Hence the mechanism for generating the density perturbations proposed in [1] is not of the kind usually considered in inflationary models where a scalar inflaton field takes non-zero values. A necessary consequence of this non-standard mechanism is nongaussian statistics for the density perturbations and the CMBR.

These consequences of the model proposed in [1] are a specific realization of the general conformal invariance considerations of our Letter. To see this we begin with the observation that a massive scalar field in de Sitter background exhibits scaling behavior for distances larger than the horizon. Indeed, its two-point function falls off with a power law [3,4]:

$$\langle \Phi(x)\Phi(x') \rangle \sim d(x, x')^{-2\gamma} \quad (1)$$

$$\gamma = \frac{3}{2} - \sqrt{\frac{9}{4} - \frac{m^2}{H^2}} \approx \frac{m^2}{3H^2}, \quad (2)$$

for $m \ll H$ and $d \gg H^{-1}$, where m is the mass of Φ and d is the geodesic distance. In spatially flat coordinates

$$d^2(x, x') = e^{H(t+t')} \left[H^{-2}(e^{-Ht} - e^{-Ht'})^2 - (\mathbf{x} - \mathbf{x}')^2 \right]. \quad (3)$$

Equation (1) is just the definition of a conformal field with dimension γ . Since the energy momentum tensor is a quadratic form in Φ involving two derivatives, it follows that the scaling dimension Δ of density perturbations due to the scalar field Φ is $\Delta = 2\gamma + 2$, in the notation of [2]. Hence the spectral index

$$n = 2\Delta - 3 = 1 + 4\gamma. \quad (4)$$

Thus for $m^2 \ll H^2$ the general formula of [2] reproduces the result of [1], namely, $n \sim 1 + \frac{4m^2}{3H^2}$.

The behavior of the two-point function (1) at large distances is a reflection of the fact that there can be no spontaneous symmetry breaking in de Sitter space [4], a situation closely analogous to the absence of spontaneous breaking of global symmetries in two dimensions. It follows that the mean value of the scalar Φ vanishes when averaging over distances $d > d_0$, where $2\gamma \ln(Hd_0) = \mathcal{O}(1)$, or for comoving wavenumbers $k \ll k_0$ with $k_0 \sim H \exp(-\frac{3H^2}{2m^2})$ as pointed out in [1]. It is this vanishing of the expectation value of the scalar field which is responsible for the fact that one must go to quadratic order in the fluctuations in Φ in order to obtain a non-vanishing result for the density perturbations in the model of [1]. This implies necessarily the existence of higher point correlations in the statistics of the density perturbations, and hence the CMBR. These higher point correlations are determined by conformal invariance as in [2], since Φ scales like a conformal field at very large distances.

Let us note the close analogy between this vanishing of the scalar field Φ when averaged over large distance scales in the model of [1] and the vanishing of the average value of the Ricci scalar due to the quantum fluctuations of the conformal factor of the spacetime metric at large distance scales [5] in the model of [2]. It is this mechanism which gives rise to conformal invariance and nongaussian statistics in the latter model. The coefficient of the trace anomaly Q^2 which gives rise to these fluctuations leads to an anomalous dimension of the scaling behavior and a “blue” spectrum for the CMBR with $4/Q^2$ playing essentially the same role as $4m^2/3H^2$. This parameter is naturally of order one in the conformal factor theory, without the need to introduce a new supersymmetric sector.

I. A. and E. M. acknowledge the hospitality of the Dept. of Physics and Astronomy, Univ. of S. Carolina.

Ignatios Antoniadis¹, Pawel O. Mazur², and Emil Mottola³

¹ Centre de Physique Théorique,

Ecole Polytechnique, 91128 Palaiseau, France and
Theory Division, CERN, 1211 Geneva 23, Switzerland

² Department of Physics and Astronomy,
University of South Carolina, Columbia, SC 29208

³ Theoretical Division, Los Alamos National Laboratory,
MS B285, Los Alamos, NM 87545

[1] A. Linde and V. Mukhanov, astro-ph/9610219.

[2] I. Antoniadis, P. O. Mazur and E. Mottola, astro-ph/9611208, to appear in Phys. Rev. Lett. (1997).

[3] N.A. Chernikov and E.A. Tagirov, Ann. Inst. Henri Poincaré **9**, 109 (1968).

[4] B. Ratra, Phys. Rev. **D31**, 1931 (1985).

[5] I. Antoniadis and E. Mottola, Phys. Rev. **D45**, 2013 (1992); I. Antoniadis, P. O. Mazur and E. Mottola, Phys. Lett. **B394**, 49 (1997); Phys. Rev. **D55**, 4756, 4770 (1997).